

QuilliChew ER

(methylphenidate ER chewable)

[Full Prescribing Information](#)

[DailyMed Drug Information](#)

Forms/Strengths

- **Chewables (Cherry-flavored):** 20 mg, 30 mg, 40 mg

Dosing

- **Age:** ≥ 6 y/o
- **Onset:** ~ 45 min
- **Duration:** up to 8 hours
- **Release Profile:** 30% IR, 70% ER
- **Considerations:** Cherry-flavored; may be taken with or without food. Long-acting chewable tablet. The 20 mg and 30 mg tablets may be split in half.
- **Initial Dose:** 20 mg
- **Titration:** 10-20 mg weekly
- **Max Dose:** 60 mg/day

Quick Facts

- Blocks dopamine/norepinephrine reuptake; enhances focus, attention, and impulse control
- Extended-release chewable tablet with dual-phase release for immediate onset and sustained effect
- Unique chewable formulation; ideal for patients with swallowing difficulties
- Common side effects: decreased appetite, insomnia, headache, gastrointestinal discomfort

Indications

- **ADHD** (ICD-10: F90.0)
-

Off-Label Uses

- **N/A**
-

How to Take

- Take **once daily in the morning** to minimize the risk of insomnia.
 - Can be taken **with or without food**; high-fat meals may delay absorption.
 - **Chew the tablet thoroughly** before swallowing—do not swallow whole.
 - Avoid taking **late in the day** to prevent sleep disturbances.
-

Side Effects

- Common: Decreased appetite, insomnia, nausea, vomiting, abdominal pain, weight loss, anxiety, dizziness, irritability, tachycardia, increased blood pressure
 - Serious: Cardiac events, psychiatric symptoms, priapism, peripheral vasculopathy, growth suppression
-

Monitoring / Labs

- **Cardiovascular:** Baseline and routine monitoring of heart rate and blood pressure.
 - **Growth in Pediatrics:** Regular monitoring of height and weight to detect growth suppression.
 - **Psychiatric Symptoms:** Observe for mood changes, anxiety, or psychosis.
 - **Abuse Potential:** Monitor for misuse or diversion.
-

Education

- **When to Call the Doctor:**
 - Severe **mood changes, aggression, or suicidal thoughts.**

- Chest pain, rapid heartbeat, shortness of breath (**cardiovascular concerns**).
 - Uncontrolled movements, tics, or worsening anxiety.
 - Numbness, coldness, or color changes in fingers or toes (**circulatory issues**).
 - Unexplained weight loss or **delayed growth in pediatric patients**.
 - **Safety Tips:**
 - **Monitor blood pressure and heart rate**, especially in patients with cardiovascular risk.
 - Use caution in patients with **anxiety, bipolar disorder, or psychosis**, as symptoms may worsen.
 - Avoid **caffeine and other stimulants**, which may amplify side effects.
 - Ensure **adequate hydration and nutrition**, as appetite suppression is common.
 - **Tapering may be required** if discontinuing after long-term use.
 - **Parent Tips for Pediatric Patients:**
 - **Encourage a nutrient-dense diet** to counteract appetite suppression.
 - Administer in the **morning before school** for optimal effect.
 - Monitor **school performance and behavioral changes**.
 - Observe for **sleep disturbances**; adjust timing if necessary.
 - Communicate regularly with **teachers and caregivers** about medication effects.
-

Additional Information

- **Contraindications:**
 - **Hypersensitivity** to methylphenidate or formulation components.
 - **Use within 14 days of MAOI therapy** (risk of hypertensive crisis).
 - Symptomatic **cardiovascular disease, moderate-to-severe hypertension, hyperthyroidism, glaucoma**.
 - **History of substance use disorder**, unless benefits outweigh risks.
 - **Pregnancy:**
 - **Category C**; use only if benefits outweigh risks.
 - May cause **neonatal withdrawal symptoms** or **low birth weight**.
 - **Lactation:**
 - **Excreted in breast milk; not recommended** due to potential infant exposure.
 - **Drug Interactions:**
 - **Serotonergic drugs** (e.g., SSRIs, SNRIs, MAOIs) increase **serotonin syndrome risk**.
 - **Acidifying agents** (e.g., ascorbic acid) may reduce drug absorption.
 - **Alkalinizing agents** (e.g., sodium bicarbonate) may increase methylphenidate levels.
 - May potentiate **hypertensive effects** of certain medications (e.g., decongestants, beta-agonists).
-

