

Quillivant XR

(methylphenidate XR suspension)

Full Prescribing Information

DailyMed Drug Information

Forms/Strengths

- **Liquid (Fruit-flavored):** 25 mg/5 mL

Dosing

- **Age:** ≥ 6 y/o
- **Onset:** ~ 45 min
- **Duration:** ~ 12 hours
- **Release Profile:** 20% IR, 80% ER
- **Considerations:** Fruit-flavored; Shake bottle vigorously for at least 10 seconds before administering. May be taken with or without food. Store at room temperature. Suspension expires four months after reconstitution.
- **Initial Dose:** 20 mg
- **Titration:** 10-20 mg weekly
- **Max Dose:** 60 mg/day

Quick Facts

- Blocks dopamine/norepinephrine reuptake; enhances focus, attention, and impulse control
- Extended-release oral suspension with dual-phase release for immediate and sustained effect
- Unique liquid formulation ideal for patients who have difficulty swallowing pills
- Common side effects: decreased appetite, insomnia, stomach upset, headache

Indications

- **ADHD** (ICD-10: F90.0)
-

Off-Label Uses

- **N/A**
-

How to Take

- Take **once daily in the morning** to minimize the risk of insomnia.
 - **Shake the bottle well** for at least **10 seconds** before each dose to ensure proper mixing.
 - **Measure the dose accurately** using the provided oral dosing syringe—do not use household spoons.
 - Can be taken **with or without food**; high-fat meals may delay absorption.
 - Avoid taking **late in the day** to prevent sleep disturbances.
-

Side Effects

- Common: Decreased appetite, insomnia, nausea, vomiting, abdominal pain, weight loss, anxiety, dizziness, irritability, tachycardia, increased blood pressure.
 - Serious: Cardiac events, psychiatric symptoms, priapism, peripheral vasculopathy, growth suppression, glaucoma, tics.
-

Monitoring / Labs

- **Cardiovascular:** Baseline and routine monitoring of heart rate and blood pressure.
 - **Growth in Pediatrics:** Regular monitoring of height and weight to detect growth suppression.
 - **Psychiatric Symptoms:** Observe for mood changes, anxiety, or psychosis.
 - **Abuse Potential:** Monitor for misuse or diversion.
-

Education

- **When to Call the Doctor:**

- Severe **mood changes, aggression, or suicidal thoughts**.
- Chest pain, rapid heartbeat, shortness of breath (**cardiovascular concerns**).
- Uncontrolled movements, tics, or worsening anxiety.
- Numbness, coldness, or color changes in fingers or toes (**circulatory issues**).
- Unexplained weight loss or **delayed growth in pediatric patients**.

- **Safety Tips:**

- **Monitor blood pressure and heart rate**, especially in patients with cardiovascular risk.
- Use caution in patients with **anxiety, bipolar disorder, or psychosis**, as symptoms may worsen.
- Avoid **caffeine and other stimulants**, which may amplify side effects.
- Ensure **adequate hydration and nutrition**, as appetite suppression is common.
- **Tapering may be required** if discontinuing after long-term use.

- **Parent Tips for Pediatric Patients:**

- **Encourage a nutrient-dense diet** to counteract appetite suppression.
- Administer in the **morning before school** for optimal effect.
- Monitor **school performance and behavioral changes**.
- Observe for **sleep disturbances**; adjust timing if necessary.
- Communicate regularly with **teachers and caregivers** about medication effects.

Additional Information

- **Contraindications:**

- **Hypersensitivity** to methylphenidate or formulation components.
- **Use within 14 days of MAOI therapy** (risk of hypertensive crisis).
- Symptomatic **cardiovascular disease, moderate-to-severe hypertension, hyperthyroidism, glaucoma**.
- **History of substance use disorder**, unless benefits outweigh risks.

- **Pregnancy:**

- **Category C**; use only if benefits outweigh risks.
- May cause **neonatal withdrawal symptoms** or **low birth weight**.

- **Lactation:**

- **Excreted in breast milk; not recommended** due to potential infant exposure.

- **Drug Interactions:**

- **Serotonergic drugs** (e.g., SSRIs, SNRIs, MAOIs) increase **serotonin syndrome risk**.
- **Acidifying agents** (e.g., ascorbic acid) may reduce drug absorption.
- **Alkalinizing agents** (e.g., sodium bicarbonate) may increase methylphenidate levels.
- May potentiate **hypertensive effects** of certain medications (e.g., decongestants, beta-agonists).

