

Medication by Form



Liquids

Short-Acting:

Methylphenidate

Methylin (methylphenidate HCl); oral solution; Brand/Generic

- **Age:** ≥ 6 y/o
- **Onset:** ~20-60 min
- **Duration:** 3-4 hours
- **Considerations:** Typically dosed multiple times per day. May be taken with or without food.
 - **Liquid:** Colorless, grape-flavored liquid; store at room temperature.
 - **Chewable:** Grape-flavored chewable tablet.
- **Initial Dose:** 5 mg twice daily (before breakfast and lunch)
- **Titration:** 5-10 mg weekly
- **Max Dose:** 2mg/kg/day up to 60 mg/day

Amphetamine

ProCentra (Dextroamphetamine sulfate); oral solution; Brand/Generic

- **Age:** ≥ 3 y/o
- **Onset:** ~ 30 min
- **Duration:** 4-6 hours
- **Considerations:** Bubblegum-flavored; may be taken with or without food. Shake bottle before administering. Store at room temperature.
- **Initial Dose:**
 - 3-5 y/o: 2.5 mg daily
 - 6+ y/o: 5 mg once or twice daily
- **Titration:**
 - 3-5 y/o: 2.5 mg weekly
 - 6+ y/o: 5 mg weekly
- **Max Dose:**
 - 3-5 y/o: 40 mg/day
 - 6+ y/o: 60 mg/day

Long-Acting:

Methylphenidate

Quillivant XR (methylphenidate XR suspension); oral suspension; Brand

- **Age:** ≥ 6 y/o
- **Onset:** ~ 45 min
- **Duration:** ~ 12 hours
- **Release Profile:** 20% IR, 80% ER
- **Considerations:** Fruit-flavored; Shake bottle vigorously for at least 10 seconds before administering. May be taken with or without food. Store at room temperature. Suspension expires four months after reconstitution.
- **Initial Dose:** 20 mg
- **Titration:** 10-20 mg weekly
- **Max Dose:** 60 mg/day

Amphetamine

Dyanavel XR (amphetamine ER); oral suspension; Brand

- **Age:** ≥ 6 y/o
- **Onset:** ~60 min
- **Duration:** up to 13 hours
- **Considerations:** May be taken with or without food. Liquid and tablet formulations are interchangeable.
 - **Tablet:** The 5mg dose is scored to allow for accurate dosing down to 2.5mg.
 - **Liquid:** Bubblegum flavor
- **Initial Dose:**
 - 6-12 y/o: 2.5-5 mg
 - 13+ y/o: 5-10 mg
- **Titration:**
 - 6-12 y/o: 2.5 mg every 4-7 days
 - 13+ y/o: 5 mg every 4-7 days
- **Max Dose:** 20 mg

Vyvanse (Lisdexamfetamine dimesylate); capsule; Brand/Generic

- **Age:** ≥ 6 y/o
- **Onset:** ~60 min
- **Duration:** 8-12 hours
- **Considerations:** Can be taken with or without food. It is a pro-drug that is metabolized into 100% dextroamphetamine, with a reduced risk of abuse. The chewable tablet and capsule are interchangeable on a mg-to-mg basis.
 - **Chewable tablet:** Strawberry-flavored and can be taken with or without food.
 - **Capsule:** The capsule can be opened and its contents dissolved in liquid, such as water, yogurt, or orange juice. Use immediately after dissolving.
- **Initial Dose:** 20-30 mg
- **Titration:** 10-20 mg every 3-7 days
- **Max Dose:** 70 mg

Chewable Tablets

Short-Acting:

Methylphenidate

Methylin (Methylphenidate HCl); chewable tablet; Generic

- **Age:** ≥ 6 y/o
- **Onset:** ~20-60 min
- **Duration:** 3-4 hours
- **Considerations:** Typically dosed multiple times per day. May be taken with or without food.
 - **Liquid:** Colorless, grape-flavored liquid; store at room temperature.
 - **Chewable:** Grape-flavored chewable tablet.
- **Initial Dose:** 5 mg twice daily (before breakfast and lunch)
- **Titration:** 5-10 mg weekly
- **Max Dose:** 2mg/kg/day up to 60 mg/day

Long-Acting:

Methylphenidate

QuilliChew ER (methylphenidate ER chewable); chewable tablet; Brand

- **Age:** ≥ 6 y/o
- **Onset:** ~ 45 min
- **Duration:** up to 8 hours
- **Release Profile:** 30% IR, 70% ER
- **Considerations:** Cherry-flavored; may be taken with or without food. Long-acting chewable tablet. The 20 mg and 30 mg tablets may be split in half.
- **Initial Dose:** 20 mg
- **Titration:** 10-20 mg weekly
- **Max Dose:** 60 mg/day

Amphetamine

Dyanavel XR (Amphetamine ER); chewable tablets; Brand

- **Age:** ≥ 6 y/o
- **Onset:** ~60 min
- **Duration:** up to 13 hours
- **Considerations:** May be taken with or without food. Liquid and tablet formulations are interchangeable.
 - **Tablet:** The 5mg dose is scored to allow for accurate dosing down to 2.5mg.
 - **Liquid:** Bubblegum flavor
- **Initial Dose:**
 - 6-12 y/o: 2.5-5 mg
 - 13+ y/o: 5-10 mg
- **Titration:**
 - 6-12 y/o: 2.5 mg every 4-7 days
 - 13+ y/o: 5 mg every 4-7 days
- **Max Dose:** 20 mg

Vyvanse (Lisdexamfetamine); chewable tablets; Brand/Generic

- **Age:** ≥ 6 y/o
- **Onset:** ~60 min
- **Duration:** 8-12 hours
- **Considerations:** Can be taken with or without food. It is a pro-drug that is metabolized into 100% dextroamphetamine, with a reduced risk of abuse. The chewable tablet and capsule are interchangeable on a mg-to-mg basis.
 - **Chewable tablet:** Strawberry-flavored and can be taken with or without food.
 - **Capsule:** The capsule can be opened and its contents dissolved in liquid, such as water, yogurt, or orange juice. Use immediately after dissolving.
- **Initial Dose:** 20-30 mg
- **Titration:** 10-20 mg every 3-7 days
- **Max Dose:** 70 mg

Oral Disintegrating Tablets (ODT)

Long-Acting:

Methylphenidate

Cotempla XR-ODT (Methylphenidate XR-ODT); ODT; Brand

- **Age:** 6-17 y/o
- **Onset:** ~ 30-60 min
- **Duration:** ~ 8 hours
- **Release Profile:** 25% IR, 75% ER
- **Considerations:** Grape-flavored, allow to dissolve in saliva. Advise to take consistently either with food or without food.
- **Initial Dose:** 17.3 mg
- **Titration:** 8.6 mg - 17.3 mg weekly
- **Max Dose:** 51.8 mg

Amphetamine

Adzenys XR-ODT (Amphetamine ER); ODT; Brand

- **Age:** ≥ 6 y/o
- **Onset:** ~ 30 min
- **Duration:** ~12 hours
- **Release Profile:** 50% IR, 50% ER
- **Considerations:** Orange-flavored. May be taken with or without food. **Allow tablet to disintegrate in saliva before swallowing.**
 - Adderall XR Equivalent Doses
- **Initial Dose:**
 - 6-17 y/o: 6.3 mg
 - 18+ y/o: 12.5 mg
- **Titration:** 3.1 mg - 6.3 mg weekly
- **Max Dose:**
 - 6-12 y/o: 18.8 mg/day
 - 13+ y/o: 12.5 mg/day

Sprinkles

Long-Acting:

Methylphenidate

Focalin XR (dexamethylphenidate XR); capsule; Brand/Generic

- **Age:** ≥ 6 y/o
- **Onset:** ~ 30 min
- **Duration:** 8-12 hours
- **Release Profile:** 50% IR, 50% ER
- **Considerations:** Capsules may be swallowed whole or opened, and the entire contents can be sprinkled on applesauce.
- **Initial Dose:**
 - **New to methylphenidate:**
 - 6-17 y/o: 5 mg
 - Adults: 10 mg
 - **Currently on methylphenidate:** Focalin XR dosage is half (1/2) the current total daily dosage of methylphenidate
 - **Currently on Focalin IR:** Give the same daily dose of Focalin XR
- **Titration:**
 - 6-17 y/o: 5 mg weekly
 - Adults: 10 mg weekly
- **Max Dose:**
 - 6-17 y/o: 30 mg/day
 - Adults: 40 mg/day

Metadate CD (methylphenidate CD); capsule; Brand

- **Age:** ≥ 6 y/o
- **Onset:** ~ 30-60 min
- **Duration:** 8-12 hours
- **Release Profile:** 30% IR, 70% ER
- **Considerations:** The capsule can be opened, and beads can be swallowed with applesauce. Take orally once daily in the morning, preferably before breakfast.
- **Initial Dose:** 20 mg
- **Titration:** 10-20 mg weekly
- **Max Dose:** 60 mg/day

Ritalin LA (methylphenidate LA); capsule; Brand/Generic

- **Age:** ≥ 6 y/o
- **Onset:** ~ 20-60 min
- **Duration:** 8-12 hours
- **Release Profile:** 50% IR, 50% ER
- **Considerations:** Capsules may be swallowed whole or opened, and the entire contents can be sprinkled on applesauce. Should not be crushed, chewed, or divided.
- **Initial Dose:** 10-20 mg
 - Patients currently using Ritalin: dosage is equivalent to Ritalin dose
- **Titration:** 10 mg weekly
- **Max Dose:** 60 mg/day

Aptensio XR (methylphenidate XR); capsule; Brand/Generic

- **Age:** ≥ 6 y/o
- **Onset:** ~ 60-120 min
- **Duration:** 8-12 hours
- **Release Profile:** 40% IR, 60% ER
- **Considerations:** Capsules may be swallowed whole or opened, and the entire contents can be sprinkled on applesauce.
- **Initial Dose:** 10 mg
- **Titration:** 10 mg weekly
- **Max Dose:** 60 mg/day

Jornay PM (methylphenidate delayed and extended release capsules); capsule; Brand

- **Age:** ≥ 6 y/o
- **Onset:** delayed until early morning (~10-12 hours after evening dose)
- **Duration:** 12-14 hours
- **Considerations:** Take in the evening between 6:30-9:30 pm. Capsules can be opened and sprinkled on applesauce.
- **Initial Dose:** 20 mg once daily in the evening
- **Titration:** 20 mg weekly
- **Max Dose:** 60 mg/day

Azstarys (Serdexmethylphenidate and dexamethylphenidate);capsule; Brand

- **Age:** ≥ 6 y/o
- **Onset:** ~ 30 min
- **Duration:** 12-14 hours
- **Release Profile:** 30% IR, 70% ER
- **Considerations:** Administer once daily in the morning, with or without food; capsules can be taken whole or opened and sprinkled onto applesauce or mixed with water.
- **Initial Dose:** 39.2/7.8 mg
- **Titration:** Increase to 52.3/10.4 mg or decrease to 26.1/5.2 mg after one week
- **Max Dose:** 52.3/10.4 mg daily

Amphetamine

Adderall XR (amphetamine salt combo ER); capsule; Brand/Generic

- **Age:** Pediatric patients ≥6 years and adults
- **Onset:** Approximately 30-60 minutes
- **Duration:** 10-12 hours
- **Release Profile:** Biphasic - an immediate-release portion followed by a delayed-release phase via beaded capsule technology
- **Considerations:** May be taken with or without food; avoid concomitant use with acidifying agents (e.g., fruit juices, vitamin C) which may decrease absorption; high-fat meals may delay T_{max} by 2.5 hours but do not affect overall bioavailability
- **Initial Dose (Pediatric):**
 - Typically 10 mg once daily in the morning
- **Titration (Pediatric):**
 - Increase by 5-10 mg increments at weekly intervals based on clinical response and tolerability
- **Max Dose (Pediatric):**
 - Generally should not exceed 30 mg per day

Amphetamine

Vyvanse (Lisdexamfetamine); capsule; Brand/Generic

- **Age:** ≥ 6 y/o
- **Onset:** ~60 min
- **Duration:** 8-12 hours
- **Considerations:** Can be taken with or without food. It is a pro-drug that is metabolized into 100% dextroamphetamine, with a reduced risk of abuse. The chewable tablet and capsule are interchangeable on a mg-to-mg basis.
 - **Chewable tablet:** Strawberry-flavored and can be taken with or without food.
 - **Capsule:** The capsule can be opened and its contents dissolved in liquid, such as water, yogurt, or orange juice. Use immediately after dissolving.
- **Initial Dose:** 20-30 mg
- **Titration:** 10-20 mg every 3-7 days
- **Max Dose:** 70 mg

Mydayis (Dextroamphetamine-Amphetamine ER Capsules); Brand/Generic

- **Age:** ≥ 13 y/o
- **Onset:** Approximately 1-2 hours
- **Duration:** Up to 16 hours
- **Considerations:** Capsules may be swallowed whole or sprinkled on cold applesauce and consumed immediately without chewing. Administer once daily upon awakening to avoid insomnia. Consistency with or without food is recommended. Do not substitute for other amphetamine products on a milligram-per-milligram basis.
- **Initial Dose:** 12.5 mg once daily in the morning
- **Titration:** 12.5 mg weekly
- **Max Dose:**
 - **13-17 y/o:** 25 mg/day
 - **Adults:** 50 mg/day

Non-Stimulant

Qelbree (viloxazine); capsule; Brand

- **Age:** ≥ 6 y/o
- **Duration:** ~ 24 hours
- **Considerations:** Bubblegum-flavored; may be taken with or without food. Shake bottle before administering. Store at room temperature.
- **Initial Dose:**
 - 6-11 y/o: 100 mg once daily
 - 12+ y/o: 200 mg once daily
- **Titration:**
 - 6-11 y/o: 100 mg weekly
 - 12+ y/o: 200 mg weekly
- **Max Dose:**
 - 6-17 y/o: 400 mg daily
 - 17+ y/o: 600 mg daily

Transdermal Patches

Long-Acting:

Methylphenidate

Daytrana (Methylphenidate patch); Brand/Generic

- **Age:** ≥ 6 y/o
- **Onset:** Approximately 2 hours after application
- **Duration:** 9-hour wear time; continuous release for 9-12 hours
- **Considerations:** Apply the patch to the hip area (alternating sites) 2 hours before the effect is needed. The patch may be removed earlier if shorter duration is desired or if late-day side effects occur. Monitor for skin rash or sensitivity. Discard patches appropriately.
- **Initial Dose:** 10 mg patch applied daily
- **Titration:** 5 mg every 3 - 7 days
- **Max Dose:** 30 mg/day

Amphetamine

Xelstrym (Dextroamphetamine patch); Brand

- **Age:** ≥ 6 y/o
- **Onset:** Approximately 2 hours after application
- **Duration:** 9-hour wear time; continuous release for 9-12 hours
- **Considerations:** Apply the patch to the hip, upper arm, chest, upper back, or flank area 2 hours before the effect is needed. Rotate application sites with each new patch. Do not use more than one patch in a 24-hour period. Do not substitute for other amphetamine products on a milligram-per-milligram basis.
- **Initial Dose:**
 - **Pediatric (6-17 y/o):** 4.5 mg/9 hours daily
 - **Adults:** 9 mg/9 hours daily
- **Titration:** 4.5 mg/9 hours weekly
- **Max Dose:** 18 mg/9 hours
 - **Severe Renal Impairment:** 13.5 mg/9 hours
 - **End-Stage Renal Disease (ESRD):** 9 mg/9 hours

Crushable Tablets

Short-Acting:

All short-acting crushable tablets listed are considered off-label when administered in this manner. While these medications are likely safe and effective when used as crushed, they

have not been FDA-approved for this specific method of administration.

Methylphenidate

Ritalin (Methylphenidate); tablet; Brand/Generic

- **Age:** ≥ 6 y/o
- **Onset:** ~ 20-60 min
- **Duration:** 3-4 hours
- **Considerations:** Abrupt onset and offset may increase the number and severity of side effects. Administer 30 to 45 minutes before meals.
- **Initial Dose:**
 - **Pediatric (6+ y/o):** 5 mg twice daily (before breakfast and lunch)
 - **Adults:** 10-15 mg twice or three times daily
- **Titration:** 5-10 mg weekly
- **Max Dose:** 60 mg/day

Focalin (Dexmethylphenidate); tablet; Brand/Generic

- **Age:** ≥ 6 y/o
- **Onset:** ~ 30 minutes
- **Duration:** 5-6 hours
- **Considerations:** Give approximately 1/2 the methylphenidate dose. Administer orally twice daily, at least 4 hours apart, with or without food.
- **Initial Dose:**
 - **New to methylphenidate:** 2.5 mg twice daily
 - **Currently taking methylphenidate:** Start with half the current total daily dose of methylphenidate (up to 10 mg twice daily)
- **Titration:** Increase by 2.5-5 mg daily every 3 to 7 days
- **Max Dose:** 20 mg/day (10 mg twice daily)

Methylin (Methylphenidate); chewable tablet; Generic

- **Age:** ≥ 6 y/o
- **Onset:** ~20-60 min
- **Duration:** 3-4 hours
- **Considerations:** Typically dosed multiple times per day. May be taken with or without food.
 - **Liquid:** Colorless, grape-flavored liquid; store at room temperature.
 - **Chewable:** Grape-flavored chewable tablet.
- **Initial Dose:** 5 mg twice daily (before breakfast and lunch)
- **Titration:** 5-10 mg weekly
- **Max Dose:** 2mg/kg/day up to 60 mg/day

Amphetamine

Adderall (amphetamine salt combo); tablets; Brand/Generic

- **Age:** ≥ 3 y/o
- **Onset:** 30-60 minutes
- **Duration:** 4-6 hours
- **Considerations:** Administer first dose upon awakening; additional doses at 4-6 hour intervals; may take with or without food (high-fat meals may delay absorption)
- **Initial Dose:**
 - **3-5 y/o:** 2.5 mg daily
 - **6+ y/o:** 5 mg 1-2 times daily
- **Titration:**
 - **3-5 y/o:** 2.5 mg weekly
 - **6+ y/o:** 5 mg weekly
- **Max Dose:**
 - **≤ 50 kg:** 40 mg/day
 - **> 50 kg:** 60 mg/day (Narcolepsy)

Zenzedi (Dextroamphetamine sulfate); tablet; Brand only at the 2.5 mg and 7.5 mg doses / Generic

- **Age:** ≥ 3 y/o
- **Onset:** ~ 30-60 minutes
- **Duration:** 4-6 hours
- **Considerations:** Take the first dose on awakening. Additional doses may be given at intervals of 4-6 hours. Periodic drug holidays are recommended to assess the patient's condition.
- **Initial Dose:**
 - **3-5 y/o:** 2.5 mg daily
 - **6+ y/o:** 5 mg once or twice daily
- **Titration:**
 - **3-5 y/o:** 2.5 mg weekly
 - **6+ y/o:** 5 mg weekly
- **Max Dose:**
 - **3-5 y/o:** 20 mg/day
 - **6+ y/o:**
 - **≤ 50 kg:** 40 mg/day
 - **> 50 kg:** 60 mg/day

Evekeo (Amphetamine sulfate); tablet; Brand

- **Age:** ≥ 3 y/o
- **Onset:** ~ 30-60 minutes
- **Duration:** 4-6 hours
- **Considerations:** Take the first dose on awakening. Additional doses may be given at intervals of 4-6 hours.
- **Initial Dose:**
 - **3-5 y/o:** 2.5 mg daily
 - **6+ y/o:** 5 mg once or twice daily
- **Titration:**
 - **3-5 y/o:** 2.5 mg weekly
 - **6+ y/o:** 5 mg weekly
- **Max Dose:** 40 mg/day

Long-Acting:

Methylphenidate

QuilliChew ER (methylphenidate ER chewable); chewable tablet; Brand

- **Age:** ≥ 6 y/o
- **Onset:** ~ 45 min
- **Duration:** up to 8 hours
- **Release Profile:** 30% IR, 70% ER
- **Considerations:** Cherry-flavored; may be taken with or without food. Long-acting chewable tablet. The 20 mg and 30 mg tablets may be split in half.
- **Initial Dose:** 20 mg
- **Titration:** 10-20 mg weekly
- **Max Dose:** 60 mg/day

Amphetamine

Dyanavel XR (Dextroamphetamine extended-release); tablet; Brand

- **Age:** ≥ 6 y/o
- **Onset:** ~60 min
- **Duration:** up to 13 hours
- **Considerations:** May be taken with or without food. Liquid and tablet formulations are interchangeable.
 - **Tablet:** The 5mg dose is scored to allow for accurate dosing down to 2.5mg.
 - **Liquid:** Bubblegum flavor
- **Initial Dose:**
 - 6-12 y/o: 2.5-5 mg
 - 13+ y/o: 5-10 mg
- **Titration:**
 - 6-12 y/o: 2.5 mg every 4-7 days
 - 13+ y/o: 5 mg every 4-7 days
- **Max Dose:** 20 mg

Vyvanse (Lisdexamfetamine); chewable tablet; Brand/Generic

- **Age:** ≥ 6 y/o
- **Onset:** ~60 min
- **Duration:** 8-12 hours
- **Considerations:** Can be taken with or without food. It is a pro-drug that is metabolized into 100% dextroamphetamine, with a reduced risk of abuse. The chewable tablet and capsule are interchangeable on a mg-to-mg basis.
 - **Chewable tablet:** Strawberry-flavored and can be taken with or without food.
 - **Capsule:** The capsule can be opened and its contents dissolved in liquid, such as water, yogurt, or orange juice. Use immediately after dissolving.
- **Initial Dose:** 20-30 mg
- **Titration:** 10-20 mg every 3-7 days
- **Max Dose:** 70 mg

Non-Chewable Tablets

Long-Acting:

Methylphenidate

Concerta (methylphenidate ER (OROS)); OSM tablets; Brand/Generic

- **Age:** \geq 6 y/o
- **Onset:** ~ 30-60 min
- **Duration:** Up to 12 hours
- **Release Profile:** 22% IR, 78% CR via osmotic delivery
- **Considerations:** Must be swallowed whole; the non-absorbable shell may pass in stool. Administer once daily in the morning, with or without food. Do not chew or crush the tablet.
- **Initial Dose:**
 - **6-12 y/o:** 18 mg daily
 - **13+ y/o:** 18 mg - 36 mg daily
- **Titration:** 18 mg/day at weekly intervals
- **Max Dose:**
 - **6-12 y/o:** 54 mg/day
 - **13+ y/o:** 72 mg/day

Relexxii (Methylphenidate ER (OROS) Tablets); Brand

Relexxii is a tablet-shaped version of Concerta, offering the same OROS delivery system and bioequivalence. It also comes in additional dosage strengths.

- **Age:** \geq 6 y/o
- **Onset:** ~60 min
- **Duration:** 8-12 hours
- **Release Profile:** 20% IR, 80% CR via osmotic delivery
- **Considerations:** Must be swallowed whole; the non-absorbable shell may pass in stool. Administer once daily in the morning, with or without food. Do not chew or crush the tablet.
- **Initial Dose:**
 - **6-12 y/o:** 18 mg once daily in the morning
 - **13+ y/o:** 18 mg once daily in the morning
- **Titration:** Increase by 18 mg/day at weekly intervals
- **Max Dose:**
 - **6-12 y/o:** 54 mg/day
 - **13+ y/o:** 72 mg/day
 - **Adults:** 72 mg/day

Non-Stimulant

Intuniv (guanfacine ER); capsule; Brand/Generic

- **Age:** ≥ 6 y/o
- **Duration:** ~ 24 hours
- **Considerations:** Avoid high-fat meals; do not crush, chew, or break tablets
- **Initial Dose:** 1 mg daily
- **Titration:** 1 mg/week
- **Max Dose:**
 - **25 - 33.9 kg:** 2 - 3 mg/day
 - **34 - 41.4 kg:** 2 - 4 mg/day
 - **41.5 - 49.4 kg:** 3 - 5 mg/day
 - **49.5 - 58.4 kg:** 3 - 6 mg/day
 - **58.5 - 91 kg:** 4 - 7 mg/day
 - **91+ kg:** 5 - 7 mg/day

Non-Stimulant

Strattera (atomoxetine); capsule; Brand/Generic

- **Age:** ≥ 6
- **Onset:** ~ 60 min
- **Considerations:** Take the first dose on awakening. Additional doses may be given at intervals of 4-6 hours.
- **Initial Dose:**
 - **≤ 70 kg:** 0.5 mg/kg/day (1 or 2 divided doses)
 - **> 70 kg:** 40 mg daily
- **Titration:**
 - **≤ 70 kg:** After at least 3 days at the initial dose, increase to approximately 1.2 mg/kg/day.
 - **> 70 kg:** After at least 3 days at 40 mg/day, increase to 80 mg/day. If no improvement after 2-4 weeks, may increase up to 100 mg/day.
- **Max Dose:**
 - **≤ 70 kg:** lesser of 1.4 mg/kg/day or 100 mg/day
 - **> 70 kg:** 100 mg/day

Revision #90

Created 21 December 2024 03:56:12 by Josh Lejeune NP

Updated 22 February 2025 14:24:26 by Josh Lejeune NP